

Une théorie locale des polylogarithmes.

Local continuation of Li.

V.C. Bùi, G.H.E. Duchamp,
Hoang Ngoc Minh, Q.H. Ngô et al.

Collaboration at various stages of the work
and in the framework of the Project

Evolution Equations in Combinatorics and Physics :
N. Behr, K. A. Penson, C. Tollu.

JNCF'19,
4-8 Février 2019

Plan

- 3 Introduction
- 4 Explicit construction of Li
- 6 Li From Noncommutative Diff.
Eq.
- 8 Properties of the extended Li
- 9 Passing to harmonic sums
 H_w , $w \in Y^*$
- 10 Global and local domains
- 11 Properties of the domains

- 15 Continuing the ladder
- 16 On the right: freeness without monodromy
- 17 A useful property
- 18 Left and then right: the arrow
 $Li_\bullet^{(1)}$
- 19 Sketch of the proof (pictorial)
- 20 Concluding remarks

Introduction

The aim of this quick talk is to explain how to extend polylogarithms

$$\text{Li}(s_1, \dots, s_r) = \sum_{n_1 > n_2 > \dots > n_r > 0} \frac{z^{n_1}}{n_1^{s_1} \dots n_r^{s_r}} \quad (1)$$

They are a priori coded by lists (s_1, \dots, s_r) but, when $s_i \in \mathbb{N}_+$, they admit an *iterated integral representation* and are better coded by words with letters in $X = \{x_0, x_1\}$. We will use the one-to-one correspondences.

$$(\textcolor{red}{s_1}, \dots, \textcolor{red}{s_r}) \in \mathbb{N}_+^r \leftrightarrow x_0^{\textcolor{red}{s_1}-1} x_1 \dots x_0^{\textcolor{red}{s_r}-1} x_1 \in X^* x_1 \leftrightarrow y_{\textcolor{red}{s_1}} \dots y_{\textcolor{red}{s_r}} \in Y^* \quad (2)$$

- $\text{Li}(s)[z]$ is Jonquière and, for $\Re(s) > 1$, one has $\text{Li}(s)[1] = \zeta(s)$
- Completed by $\text{Li}(x_0^n) = \frac{\log^n(z)}{n!}$ this provides a family of independant functions admitting an analytic continuation on the cleft plane $\mathbb{C} \setminus (]-\infty, 0] \cup [1, +\infty[)$ or $\widetilde{\mathbb{C} \setminus \{0, 1\}}$.

Explicit construction of Li


Given a word w , we note $|w|_{x_1}$ the number of occurrences of x_1 within w

$$\alpha_0^z(w) = \begin{cases} 1_\Omega & \text{if } w = 1x^* \\ \int_0^z \alpha_0^s(u) \frac{ds}{1-s} & \text{if } w = x_1 u \\ \int_1^z \alpha_0^s(u) \frac{ds}{s} & \text{if } w = x_0 u \text{ and } |u|_{x_1} = 0 \\ \int_0^z \alpha_0^s(u) \frac{ds}{s} & \text{if } w = x_0 u \text{ and } |u|_{x_1} > 0 . \end{cases} \quad (3)$$

Of course, the third line of this recursion implies

$$\alpha_0^z(x_0^n) = \frac{\log(z)^n}{n!}$$

one can check that (a) all the integrals (although improper for the fourth line) are well defined (b) the series $S = \sum_{w \in X^*} \alpha_0^z(w) w$ satisfies (4). We then have $\alpha_0^z = \text{Li}$.



As an example, we compute some coefficients

$$\langle \text{Li} | x_0^n \rangle = \frac{\log(z)^n}{n!} \quad ; \quad \langle \text{Li} | x_1^n \rangle = \frac{(-\log(1-z))^n}{n!}$$

$$\langle \text{Li} | x_0x_1 \rangle = \text{Li}_2(z) = \sum_{n \geq 1} \frac{z^n}{n^2} \quad ; \quad \langle \text{Li} | x_1x_0 \rangle = \langle \text{Li} | x_1 \sqcup x_0 - x_0x_1 \rangle(z)$$

$$\langle \text{Li} | x_0^2x_1 \rangle = \text{Li}_3(z) = \sum_{n \geq 1} \frac{z^n}{n^3} \quad ; \quad \langle \text{Li} | x_1x_0 \rangle = (-\log(1-z))\log(z) - \text{Li}_2(z)$$

$$\langle \text{Li} | x_0^{r-1}x_1 \rangle = \text{Li}_r(z) = \sum_{n \geq 1} \frac{z^n}{n^r} \quad ; \quad \langle \text{Li} | x_1^2x_0 \rangle = \langle \text{Li} | \frac{1}{2}(x_1 \sqcup x_1 \sqcup x_0) - (x_1 \sqcup x_0x_1) + x_0x_1^2 \rangle$$

Li From Noncommutative Diff. Eq.

The generating series $S = \sum_{w \in X^*} Li(w)$ satisfies (and is unique to do so)

$$\left\{ \begin{array}{l} \mathbf{d}(S) = \left(\frac{x_0}{z} + \frac{x_1}{1-z} \right) \cdot S \\ \lim_{\substack{z \rightarrow 0 \\ z \in \Omega}} S(z) e^{-x_0 \log(z)} = 1_{\mathcal{H}(\Omega) \langle\langle X \rangle\rangle} \end{array} \right. \quad (4)$$

with $X = \{x_0, x_1\}$. This is, up to the sign of x_1 , the solution G_0 of Drinfel'd [2] for KZ3. We define this unique solution as Li . All Li_w are \mathbb{C} - and even $\mathbb{C}(z)$ -linearly independant (see CAP 17 *Linear independance without monodromy* [5]).

Domain of Li (definition)

In order to extend indexation of Li to series, we define $\text{Dom}(Li; \Omega)$ (or $\text{Dom}(Li)$) if the context is clear) as the set of series $S = \sum_{n \geq 0} S_n$ (decomposition by homogeneous components) such that $\sum_{n \geq 0} Li_{S_n}(z)$ converges unconditionally for compact convergence in Ω . One sets

$$Li_S(z) := \sum_{n \geq 0} Li_{S_n}(z) \quad (5)$$

Starting the ladder

$$\begin{array}{ccc} (\mathbb{C}\langle X \rangle, \sqcup, 1_{X^*}) & \xrightarrow{\text{Li}_\bullet} & \mathbb{C}\{\text{Li}_w\}_{w \in X^*} \\ \downarrow & & \downarrow \\ (\mathbb{C}\langle X \rangle, \sqcup, 1_{X^*})[x_0^*, (-x_0)^*, x_1^*] & \xrightarrow{\text{Li}_\bullet^{(1)}} & \mathcal{C}_{\mathbb{Z}}\{\text{Li}_w\}_{w \in X^*} \end{array}$$

Examples

$$Li_{x_0^*}(z) = z, \quad Li_{x_1^*}(z) = (1 - z)^{-1}, \quad Li_{\alpha x_0^* + \beta x_1^*}(z) = z^\alpha (1 - z)^{-\beta}$$

Properties of the extended Li

Proposition

With this definition, we have

- ① $\text{Dom}(Li)$ is a shuffle subalgebra of $\mathbb{C}\langle\langle X \rangle\rangle$ and so is

$$\text{Dom}^{\text{rat}}(Li) := \text{Dom}(Li) \cap \mathbb{C}^{\text{rat}}\langle\langle X \rangle\rangle$$

- ② For $S, T \in \text{Dom}(Li)$, we have

$$\text{Li}_{S \sqcup T} = \text{Li}_S \cdot \text{Li}_T$$

Examples and counterexamples

For $|t| < 1$, one has $(tx_0)^*x_1 \in \text{Dom}(Li, D)$ (D being the open unit slit disc and $\text{Dom}(Li, D)$ defined similarly), whereas $x_0^*x_1 \notin \text{Dom}(Li, D)$.

Indeed, we have to examine the convergence of $\sum_{n \geq 0} \text{Li}_{x_0^n x_1}(z)$, but, for $z \in]0, 1[$, one has $0 < z < \text{Li}_{x_0^n x_1}(z) \in \mathbb{R}$ and therefore, for these values $\sum_{n \geq 0} \text{Li}_{x_0^n x_1}(z) = +\infty$. Contrariwise one can show that, for $|t| < 1$,

$$\text{Li}_{(tx_0)^*x_1}(z) = \sum_{n \geq 1} \frac{z^n}{n-t}$$

Passing to harmonic sums H_w , $w \in Y^*$

Polylogarithms having a removable singularity at zero

The following proposition helps us characterize their indices.

Proposition

Let $f(z) = \langle L \mid P \rangle = \sum_{w \in X^*} \langle P \mid w \rangle \text{Li}_w$. The following conditions are equivalent

- i) f can be analytically extended around zero
- ii) $P \in \mathbb{C}\langle X \rangle x_1 \oplus \mathbb{C}.1_{X^*}$

We recall the expansion (for $w \in X^* x_1 \sqcup \{1_{X^*}\}$, $|z| < 1$)

$$\frac{\text{Li}_w(z)}{1-z} = \sum_{N \geq 0} H_{\pi_Y(w)}(N) z^N \quad (6)$$

Global and local domains

This proposition and the lemma lead us to the following definitions.

① Global domains.–

Let $\emptyset \neq \Omega \subset \widetilde{B}$ (with $B = \mathbb{C} \setminus \{0, 1\}$), we define $Dom_{\Omega}(Li) \subset \mathbb{C}\langle\langle X \rangle\rangle$ to be the set of series $S = \sum_{n \geq 0} S_n$ (with $S_n = \sum_{|w|=n} \langle S \mid w \rangle w$ each homogeneous component) such that $\sum_{n \in \mathbb{N}} Li_{S_n}$ is unconditionally convergent for the compact convergence (UCC) [4].

As examples, we have Ω_1 , the doubly cleft plane then

$$Dom(Li) := Dom_{\Omega_1}(Li) \text{ or } \Omega_2 = \widetilde{B}$$

② Local domains around zero (fit with H-theory).–

Here, we consider series $S \in (\mathbb{C}\langle\langle X \rangle\rangle x_1 \oplus \mathbb{C} 1_{X^*})$ (i.e. $supp(S) \cap Xx_0 = \emptyset$).

We consider radii $0 < R \leq 1$, the corresponding open discs

$$D_R = \{z \in \mathbb{C} \mid |z| < R\} \text{ and define}$$

$$Dom_R(Li) := \{S = \sum_{n \geq 0} S_n \in (\mathbb{C}\langle\langle X \rangle\rangle x_1 \oplus \mathbb{C} 1_{\Omega}) \mid \sum_{n \in \mathbb{N}} Li_{S_n} \text{ (UCC) in } D_R\}$$

$$Dom_{loc}(Li) := \cup_{0 < R \leq 1} Dom_R(Li).$$

Properties of the domains

Theorem A

- ① For all $\emptyset \neq \Omega \subset \widetilde{B}$, $Dom_{\Omega}(\text{Li})$ is a shuffle subalgebra of $\mathbb{C}\langle\langle X \rangle\rangle$ and so are the $Dom_R(\text{Li})$.
- ② $R \mapsto Dom_R(\text{Li})$ is strictly decreasing for $R \in]0, 1]$.
- ③ All $Dom_R(\text{Li})$ and $Dom_{loc}(\text{Li})$ are shuffle subalgebras of $\mathbb{C}\langle\langle X \rangle\rangle$ and $\pi_Y(Dom_{loc}(\text{Li}))$ is a stuffle subalgebra of $\mathbb{C}\langle\langle Y \rangle\rangle$.
- ④ Let $T(z) = \sum_{N \geq 0} a_N z^N$ be a Taylor series i.e. such that $\limsup_{N \rightarrow +\infty} |a_N|^{1/N} = B < +\infty$, then the series

$$S = \sum_{N \geq 0} a_N (-(-x_1)^+)^{\sqcup N} \quad (7)$$

is summable in $\mathbb{C}\langle\langle X \rangle\rangle$ (with sum in $\mathbb{C}\langle\langle x_1 \rangle\rangle$) and $S \in Dom_R(\text{Li})$ with $R = \frac{1}{B+1}$ and $\text{Li}_S = T(z)$.

Theorem A/2

⑤ Let $S \in Dom_R(\text{Li})$ and $S = \sum_{n \geq 0} S_n$ (homogeneous decomposition), we define^a $N \mapsto H_{\pi_Y(S)}(N)$ by

$$\frac{\text{Li}_S(z)}{1-z} = \sum_{N \geq 0} H_{\pi_Y(S)}(N) z^N. \quad (8)$$

Moreover, for all $r \in]0, R[$, we have

$$\sum_{n, N \geq 0} |H_{\pi_Y(S_n)}(N)| r^N < +\infty, \quad (9)$$

in particular, for all $N \in \mathbb{N}$ the series (of complex numbers) $\sum_{n \geq 0} H_{\pi_Y(S_n)}(N)$ converges absolutely to $H_{\pi_Y(S)}(N)$.

^aThis definition is compatible with the old one when S is a polynomial.

Theorem A/3

- ⑥ Conversely, let $Q \in \mathbb{C}\langle\langle Y \rangle\rangle$ with $Q = \sum_{n \geq 0} Q_n$ (decomposition by weights), we suppose that it exists $r \in]0, 1]$ such that

$$\sum_{n, N \geq 0} |\mathcal{H}_{Q_n}(N) r^N| < +\infty \quad (10)$$

in particular, for all $N \in \mathbb{N}$, $\sum_{n \geq 0} \mathcal{H}_{Q_n}(N) = \ell(N) \in \mathbb{C}$ unconditionally.

Under such circumstances, $\pi_X(Q) \in Dom_r(\text{Li})$ and, for all $|z| \leq r$

$$\frac{\text{Li}_S(z)}{1-z} = \sum_{N \geq 0} \ell(N) z^N, \quad (11)$$




Figure: Jacques Hadamard and Paul Montel.

Continuing the ladder

$$\begin{array}{ccc} (\mathbb{C}\langle X \rangle, \sqcup, 1_{X^*}) & \xrightarrow{\text{Li}_\bullet} & \mathbb{C}\{\text{Li}_w\}_{w \in X^*} \\ \downarrow & & \downarrow \\ (\mathbb{C}\langle X \rangle, \sqcup, 1_{X^*})[x_0^*, (-x_0)^*, x_1^*] & \xrightarrow{\text{Li}_\bullet^{(1)}} & \mathcal{C}_{\mathbb{Z}}\{\text{Li}_w\}_{w \in X^*} \\ \downarrow & & \downarrow \\ \mathbb{C}\langle X \rangle \sqcup \mathbb{C}^{\text{rat}}\langle\langle x_0 \rangle\rangle \sqcup \mathbb{C}^{\text{rat}}\langle\langle x_1 \rangle\rangle & \xrightarrow{\text{Li}_\bullet^{(2)}} & \mathcal{C}_{\mathbb{C}}\{\text{Li}_w\}_{w \in X^*} \\ \uparrow & \nearrow & \downarrow \\ \mathbb{C}\langle X \rangle \otimes_{\mathbb{C}} \mathbb{C}^{\text{rat}}\langle\langle x_0 \rangle\rangle \otimes_{\mathbb{C}} \mathbb{C}^{\text{rat}}\langle\langle x_1 \rangle\rangle & & \end{array}$$

We have, after a theorem by Leopold Kronecker,

$$\mathbb{C}^{\text{rat}}\langle\langle x \rangle\rangle = \left\{ \frac{P}{Q} \right\}_{\substack{P, Q \in \mathbb{C}[x] \\ Q(0) \neq 0}} \quad (12)$$

On the right: freeness without monodromy

Theorem (Deneufchâtel, GHED, Minh & Solomon, 2011 [1])

Let (\mathcal{A}, ∂) be a k -commutative associative differential algebra with unit and \mathcal{C} be a differential subfield of \mathcal{A} (i.e. $\partial(\mathcal{C}) \subset \mathcal{C}$). We suppose that $k = \ker(\partial)$ and that $S \in \mathcal{A}\langle\langle X \rangle\rangle$ is a solution of the differential equation

$$\mathbf{d}(S) = MS ; \quad \langle S | 1 \rangle = 1 \text{ with } M = \sum_{x \in X} u_x \color{red}{x} \in \mathcal{C}\langle\langle X \rangle\rangle \quad (13)$$

(i.e. M is a homogeneous series of degree 1)

The following conditions are equivalent :

- ① The family $(\langle S | w \rangle)_{w \in X^*}$ of coefficients of S is (linearly) free over \mathcal{C} .
- ② The family of coefficients $(\langle S | x \rangle)_{x \in X \cup \{1_{X^*}\}}$ is (linearly) free over \mathcal{C} .
- ③ The family $(u_x)_{x \in X}$ is such that, for $f \in \mathcal{C}$ et $\alpha_x \in k$

$$\partial(f) = \sum_{x \in X} \alpha_x u_x \implies (\forall x \in X)(\alpha_x = 0).$$

A useful property

Independence of characters with respect to polynomials

▲ I came across the following property :

5 Let \mathfrak{g} be a Lie algebra over a ring k without zero divisors, $\mathcal{U} = \mathcal{U}(\mathfrak{g})$ be its enveloping algebra. As such, \mathcal{U} is a Hopf algebra and ϵ , its counit, is the only character of $\mathcal{U} \rightarrow k$ which vanishes on \mathfrak{g} .

▼ Set $\mathcal{U}_+ = \ker(\epsilon)$. We build the following filtrations ($N \geq 1$)

1

$$\mathcal{U}_N = \mathcal{U}_+^N = \underbrace{\mathcal{U}_+ \dots \mathcal{U}_+}_{N \text{ times}} \quad (1)$$

and

$$\mathcal{U}_N^* = \mathcal{U}_{N+1}^\perp = \{f \in \mathcal{U}^* \mid (\forall u \in \mathcal{U}_{N+1})(f(u) = 0)\} \quad (2)$$

the first one is decreasing and the second one increasing. One shows easily that (with \diamond as the convolution product)

$$\mathcal{U}_p^* \diamond \mathcal{U}_q^* \subset \mathcal{U}_{p+q}^*$$

so that $\mathcal{U}_\infty^* = \cup_{n \geq 1} \mathcal{U}_n^*$ is a convolution subalgebra of \mathcal{U}^* .

Now, we can state the

Theorem : The set of characters of $(\mathcal{U}, \cdot, 1_{\mathcal{U}})$ is linearly free w.r.t. \mathcal{U}_∞^* .

asked 1 month ago
viewed 106 times

FEATURED ON META

- Revisiting the "Hot Network Questions" feature, what are our shared goals for...
- Who cut the cheese?
- Responsive design released for all Beta & Undesigned sites

Related

- 39 What does the generating function $x/(1 - e^{-x})$ count?
- 11 Is there a canonical Hopf structure on the center of a universal enveloping algebra?
- 4 Do stunted exponential series give projections of a cocommutative bialgebra on its coradical filtration?
- 27 quantum groups... not via presentations
- 3 How a unitary corepresentation of a Hopf C*-algebra, deals with the antipode?
- 14 Rui Almeida with Hopf restricted (or Sweedler)

Left and then right: the arrow $\text{Li}_\bullet^{(1)}$

Proposition

- i. The family $\{x_0^*, x_1^*\}$ is algebraically independent over $(\mathbb{C}\langle X \rangle, \sqcup, 1_{X^*})$ within $(\mathbb{C}\langle\langle X \rangle\rangle^{\text{rat}}, \sqcup, 1_{X^*})$.
- ii. $(\mathbb{C}\langle X \rangle, \sqcup, 1_{X^*})[x_0^*, x_1^*, (-x_0)^*]$ is a free module over $\mathbb{C}\langle X \rangle$, the family $\{(x_0^*)^{\sqcup k} \sqcup (x_1^*)^{\sqcup l}\}_{(k,l) \in \mathbb{Z} \times \mathbb{N}}$ is a $\mathbb{C}\langle X \rangle$ -basis of it.
- iii. As a consequence, $\{w \sqcup (x_0^*)^{\sqcup k} \sqcup (x_1^*)^{\sqcup l}\}_{\substack{w \in X^* \\ (k,l) \in \mathbb{Z} \times \mathbb{N}}}$ is a \mathbb{C} -basis of it.
- iv. $\text{Li}_\bullet^{(1)}$ is the unique morphism from $(\mathbb{C}\langle X \rangle, \sqcup, 1_{X^*})[x_0^*, (-x_0)^*, x_1^*]$ to $\mathcal{H}(\Omega)$ such that

$$x_0^* \rightarrow z, \quad (-x_0)^* \rightarrow z^{-1} \text{ and } x_1^* \rightarrow (1-z)^{-1}$$

- v. $\text{Im}(\text{Li}_\bullet^{(1)}) = \mathcal{C}_{\mathbb{Z}}\{\text{Li}_w\}_{w \in X^*}$.
- vi. $\ker(\text{Li}_\bullet^{(1)})$ is the (shuffle) ideal generated by $x_0^* \sqcup x_1^* - x_1^* + 1_{X^*}$.

Sketch of the proof (pictorial)




Figure: Rewriting mod \mathcal{J} of $\{w \sqcup (x_0^*)^{\sqcup k} \sqcup (x_1^*)^{\sqcup l}\}_{k \in \mathbb{Z}, l \in \mathbb{N}, w \in X^*}$.

Concluding remarks

- Extending the domain of polylogarithms to (some) rational series permits the projection of rational identities. Such as

$$(\alpha x)^* \sqcup (\beta y)^* = (\alpha x + \beta y)^*$$

- The theory developed here allows to pursue, for the Harmonic sums, this investigation such as

$$(\alpha y_i)^* \sqcup (\beta y_j)^* = (\alpha y_i + \beta y_j + \alpha\beta y_{i+j})^*$$

- More in Minh's talk.

- [1] M. Deneufchâtel, GHED, Hoang Ngoc Minh, A. I. Solomon.–
Independence of hyperlogarithms over function fields via algebraic combinatorics, Lecture Notes in Computer Science (2011), Volume 6742 (2011), 127-139.
- [2] V.G. Drinfel'd, *On quasitriangular quasi-hopf algebra and a group closely connected with $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$* , Leningrad Math. J., 4, 829-860, 1991.
- [3] J. Hadamard, *Théorème sur les séries entières*, Acta Math., Vol 22 (1899), 55-63.
- [4] P. Montel.– *Leçons sur les familles normales de fonctions analytiques et leurs applications*, Gauthier-Villars (1927)
- [5]
<https://lipn.univ-paris13.fr/~ngocminh/CAP3.html>




Figure: ... and a lot of (machine) computations.

THANK YOU FOR YOUR ATTENTION !

Combinatorics and Arithmetic for Physics: special days



24-25 October 2018

Amphithéâtre Léon Motchane

Bures-sur-Yvette, France

Organisers

Gérard H. E. DUCHAMP
Maxim KONTSEVITCH
Gleb KOSHEVOY
Hoang Ngoc MINH

Speakers

Nicolas Behr (IRIF Univ. Paris Diderot)
Alin Bostan (INRIA Saclay)
Marek Bozejko (Univ. of Wroclaw)
Pierre Cartier (IHES)
Gérard H.E. Duchamp (IHP & Paris 13)
Dimitri Grigoryev (CNRS-Lille 1)
Dmitry Gurevich (Univ. Valenciennes)
Natalia Iyudu (Univ. Edinburgh & IHES)
Richard Kerner (LPTMC)
Gleb Koshevoy (Poncelet Lab, Moscow)
Hoàng Ngoc Minh (Lille 2 & LIPN)
Karol Penson (LPTMC & Paris 6)
Pierre Vanhove (CEA/Saclay)

Sponsor

GDR

« Renormalisation »
renorm.math.cnrs.fr

Web announcement

lipn.univ-paris13.fr/~duchamp/Conferences/CAPS_v3.html



ANNALES DE L'INSTITUT HENRI POINCARÉ D

Combinatorics, Physics and their Interactions

[Current Issue](#) | [All Issues](#) | [Author Index](#) | [Title Index](#) | [Online-First](#) | [Forthcoming](#)

[Editorial Board](#) | [Subscriptions](#) | [Instructions for Authors](#) | [Instructions for Editors](#)

The unfolding of new ideas in physics is often tied to the development of new combinatorial methods, and conversely some problems in combinatorics have been successfully attacked using methods inspired by statistical physics or quantum field theory.

The journal is dedicated to publishing high-quality original research articles and survey articles in which combinatorics and physics interact in both directions. Combinatorial papers should be motivated by potential applications to physical phenomena or models, while physics papers should contain some interesting combinatorial development.

Both rigorous mathematical proof and heuristic physical reasoning have a place in this journal, but each must be clearly labeled. Definitions and proofs should be presented with the precision and rigor that are expected in a mathematics journal. Conjectures based on heuristic physical arguments and/or numerical evidence are warmly encouraged, but they should be clearly labeled as such and should be stated as precisely as possible.

Authors should remember that readers of the journal will come from a wide variety of backgrounds, both mathematical (not only combinatorics but also algebra, topology, geometry, analysis, probability, etc) and physical (quantum field theory, statistical mechanics, quantum gravity etc). Therefore, authors should write with such a diverse audience in mind and should take care to provide, in the introductory section of their article, a clear statement of the problem to be studied and the mathematical and/or physical background from which it arises. Authors are also encouraged to provide, in the second section of the article, a brief pedagogical review of the mathematical or physical tools that will be used in their work.

The list of specific subject areas in which articles are anticipated includes:

- Combinatorics of renormalization
- Combinatorics of cluster, virial and related expansions
- Discrete geometry and combinatorics of quantum gravity
- Graph polynomials and statistical-mechanics models
- Topological graph polynomials and quantum field theory
- Physical applications of combinatorial Hopf algebras, matroids, combinatorial species, and other combinatorial structures
- Exact solutions of statistical-mechanical models
- Combinatorics and algebra of integrable systems
- Computational complexity and its relation with statistical physics
- Computational/algorithmic aspects of combinatorial physics
- Interactions of combinatorial physics with topology, geometry, probability theory, or computer science

DOI: 10.4171/AIHPD



Figure: CAP & AIHP-D.

JNCF'19, 4-8 Février 2019

V.C. Bùi, G.H.E. Duchamp, Hoang Ngoc Min

Une théorie locale des polylogarithmes.

24 / 24